Neuroscience Letters, 74 (1987) 383–387
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abens, J., see Westlind-Danielsson, A., 237
Alstermark, B., Kümmel, H. and Tantisira, B., Monosynaptic raphspinal and reticulospinal projection to forelimb motoneurones in cats, 286
Alstermark, B., Kümmel, H., Pinter, M.J. and Tantisira, B., Branching and termination of C_3–C_4 propriospinal neurones in the cervical spinal cord of the cat, 291
Amato, G., see Cortimiglia, R., 193
Ammann, D., Chao, P. and Simon, W., Valinomycin-based K^+-selective microelectrodes with low electrical membrane resistance, 221
Andell, S., see Westlind-Danielsson, A., 237
Andersen, A.C., see Cailliez, D., 163
Assenmacher, I., see Ixart, G., 85
Astier, B., Kitahama, K., Denoroy, L., Jouvet, M. and Renaud, B., Immunohistochemical evidence for the adrenergic medullary longitudinal bundle as a major ascending pathway to the locus coeruleus, 132
Aulisi, E.F., see Valentino, R.J., 297

Bandler, R. and Tork, I., Midbrain periaqueductal grey region in the cat has afferent and efferent connections with solitary tract nuclei, 1
Barbanel, G., see Ixart, G., 85
Bardasano, J.L., see Cos, S., 119
Bartfai, T., see Westlind-Danielsson, A., 237
Beal, J.A. and Knight, D.S., Classification of aberrant primary afferents in the substantia gelatinosa of the rat following neonatal capsaicin treatment, 139
Biliotti, G., see Ferri, G.-L., 304
Björklund, H., see Hermanson, A., 377
Blanchard, J.-C., see Böhme, G.A., 227
Bockaert, J., see Recasens, M., 211
Böhme, G.A., Stutzmann, J.-M., Roques, B.P. and Blanchard, J.-C., Effects of selective μ- and δ-opioid peptides on kindled amygdaloid seizures in rats, 227
Bowers, M., see Ross, C.D., 205
Brailowsky, S., Menini, C., Silva-Barrat, C. and Naquet, R., Epileptogenic γ-aminobutyric acid-withdrawal syndrome after chronic, intracortical infusion in baboons, 75
Brini, A., see Panerai, A.E., 348
Buckley, N.J., see Hassall, C.J.S., 145
Bugnon, C., see Fellmann, D., 275
Buma, P. and Nieuwenhuys, R., Ultrastructural demonstration of oxytocin and vasopressin release sites in the neural lobe and median eminence of the rat by tannic acid and immunogold methods, 151
Burnstock, G., see Hassall, C.J.S., 145

Cailliez, D., Danger, J.-M., Polak, J.M., Pelletier, G., Andersen, A.C., Leboulenger, F. and Vaudry, H., Co-distribution of neuropeptide Y and its C-terminal flanking peptide in the brain and pituitary of the frog *Rana ridibunda*, 163
Carlstedt, T., Dalsgaard, C.-J. and Molander, C., Regrowth of lesioned dorsal root nerve fibers into the spinal cord of neonatal rats, 14
Carmody, J. and Cooper, K., Swim stress reduces chronic pain in mice through an opioid mechanism, 358
Chao, P., see Ammann, D., 221
Chu, D.C.M., see Maragos, W.F., 371
Cohen, G., see Slivka, A., 112
Conte-Devolx, B., see Ixart, G., 85
Cooper, K., see Carmody, J., 358
Cortimiglia, R., Crescimanno, G., Salerno, M.T., Amoto, G. and Infantellina, F., Electrophysiological relationship between claustrum and contralateral area 4 and 6 pyramidal tract neurons, 193
Cos, S., Bardasano, J.L., Mediavilla, M.D. and Sánchez Barceló, E.J., Myeloid bodies associated with lipid droplets in pinealocytes of blind, blind-bulbectomized, blind-underfed or blind cold-exposed rats, 119
Crescimanno, G., see Cortimiglia, R., 193
Croughs, R.J.M., see Stolp, R., 337

Daigle, M., see Di Paolo, T., 250
Dalsgaard, C.-J., see Carlstedt, T., 14
Dalsgaard, C.-J., see Hermanson, A., 377
Dalsgaard, C.-J., see Molander, C., 37
D'Amato, C.J., see Maragos, W.F., 371
Danger, J.-M., see Cailliez, D., 163
Denoroy, L., see Astier, B., 132
Deutch, A.Y. and Roth, R.H., Calcitonin gene-related peptide in the ventral tegmental area: se-

0304-3940/87/$ 03.50 © 1987 Elsevier Scientific Publishers Ireland Ltd.

lective modulation of prefrontal cortical dopamine metabolism, 169
Di Paolo, T., Harnois, C. and Daigle, M., Assay of dopamine and its metabolites in human and rat retina, 250
Di Paolo, T., see Falardeau, P., 43
Dockray, G.J., see Ferri, G.-L., 304
Dosemeci, A. and Rodnight, R., Demonstration by phase-partitioning in Triton X-114 solutions that phosphoprotein B-50 (F-1) from rat brain is an integral membrane protein, 325
Dragunow, M., Time-dependent effects of papaverine on electrically induced seizures in rat, 320
Dubois, A., see Savasta, M., 180

Falardeau, P. and Di Paolo, T., Regional effect of estradiol on rat caudate-putamen dopamine receptors: lateral–medial differences, 43
Fasmer, O.B., Post, C. and Hole, K., Increased sensitivity to intrathecal substance P following chronic administration of zimelidine, 81
Fellmann, D., Bugnon, C. and Risold, P.Y., Unrelated peptide immunoreactivities coexist in neurons of the rat lateral dorsal hypothalamus: human growth hormone-releasing factor$_{1-37}$, salmon melanin-concentrating hormone- and α-melanotropin-like substances, 275
Ferri, G.-L., Morreale, R.A., Soimero, L., Biliotti, G. and Dockray, G.J., Intramural distribution of Met^5-enkephalin-Arg^6-Gly^7-Leu^8 in sphincter regions of the human gut, 304
Feuerstein, C., see Savasta, M., 180
Fujioka, T., see Ohmori, Y., 175

German, D.C., see Saper, C.B., 364
Godfrey, D.A., see Ross, C.D., 205
Goldfinger, M.D., see Pearson, J.C., 125
Grenon, G., see Robitaille, R., 187
Grino, M., see Ixart, G., 85

Häggblad, J. and Heilbronn, E., Externally applied adenosine-5′-triphosphate causes inositol triphosphate accumulation in cultured chick myotubes, 199
Hallanger, A.E., see Levey, A.I., 7
Hansen, S., see Perry, T.L., 217
Hari, R., see Huttunen, J., 63
Harnois, C., see Di Paolo, T., 250
Hasegawa, S., see Naoi, M., 232
Hassall, C.J.S., Buckley, N.J. and Burnstock, G., Autoradiographic localisation of muscarinic receptors on guinea pig intracardiac neurones and atrial myocytes in culture, 145
Hayling, N., see Ward, L.C., 353
Heilbronn, E., see Häggblad, J., 199
Henriksen, S.J., see Wiesner, J.B., 95
Hermanson, A., Dalsgaard, C.-J., Björklund, H. and Lindblom, U., Sensory reinnervation and sensibility after superficial skin wounds in human patients, 377
Hole, K., see Fasmer, O.B., 81
Holstege, G., Anatomical evidence for an ipsilateral rubrospinal pathway and for direct rubrospinal projections to motoneurons in the cat, 269
Huttunen, J. and Hari, R., Long-latency OFF-responses from the human sensorimotor cortex to tetanizing stimulation of thenar muscles, 63

Igano, K., see Itoh, H., 102
Imura, H., see Itoh, H., 102
Infantellina, F., see Cortimiglia, R., 193
Inouye, K., see Itoh, H., 102
Ishiki, R., see Naoi, M., 232
Itoh, H., Nakao, K., Katsuura, G., Morii, N., Shiono, S., Yamada, T., Sugawara, A., Saito, Y., Watanabe, K., Igano, K., Inouye, K. and Imura, H., Atrial natriuretic polypeptides: structure–activity relationship in the central action – a comparison of their antidipsogenic actions, 102
Ixart, G., Barbanel, G., Conte-Devolx, B., Grino, M., Oliver, C. and Assenmacher, I., Evidence for basal and stress-induced release of corticotropin releasing factor in the push–pull cannulated median eminence of conscious free-moving rats, 85

Jones, K., see Perry, T.L., 217
Jones, S.W., A muscarine-resistant M-current in C cells of bullfrog sympathetic ganglia, 309
Jouvet, M., see Astier, B., 132

Kaila, K., Rydqvist, B., Swerup, C. and Voipio, J., Stimulation-induced changes in the intracellular sodium activity of the crayfish stretch receptor, 53
Katsuura, G., see Itoh, H., 102
Kavaliers, M., Aggression and defeat-induced opioid analgesia displayed by mice are modified by calcium channel antagonists and agonists, 107
Kennedy, P.R., Light labeling of red nucleus neurons following an injection of peroxidase-conjugated wheat germ agglutinin into the inferior olivary nucleus of the rat, 262

Kihara, M., see Kubo, T., 331
Király, I. and van Ree, J.M., Behavioral evidence for the involvement of endogenous opioids in the interaction between cholecystokinin and brain dopamine systems, 343
Kitahama, K., see Astier, B., 132
Knight, D.S., see Beal, J.A., 139
Krnjević, K., see Zhang, L., 58
Kubo, T. and Kihara, M., Evidence for the presence of GABAergic and glycine-like systems responsible for cardiovascular control in the nucleus tractus solitarii of the rat, 331
Kümmel, H., see Alstermark, B., 286
Kümmel, H., see Alstermark, B., 291

Leboulenger, F., see Cailliez, D., 163
Levey, A.I., Hallanger, A.E. and Wainer, B.H., Cholinergic nucleus basalis neurons may influence the cortex via the thalamus, 7
Lindblom, U., see Hermanson, A., 377
Lindefors, N., Tossman, U. and Ungerstedt, U., Subchronic haloperidol and sulpiride treatment induces region-specific changes in tissue levels of putative amino acid transmitters in rat brain, 90
Liu, K.P., see Tamir, H., 257

Manier, M., see Savasta, M., 180
Mantegazza, P., see Panerai, A.E., 348
Maragos, W.F., Chu, D.C.M., Young, A.B., D'Amato, C.J. and Penney, Jr., J.B., Loss of hippocampal [^{3}H]TCP binding in Alzheimer's disease, 371
Martinez, S., see Martinez-de-la-Torre, M., 31
Martinez-de-la-Torre, M., Martinez, S. and Puelles, L., Solitary magnocellular neurons in the avian optic tectum: cytoarchitectonic, histochemical and [^{3}H]thymidine autoradiographic characterization, 31
Mason, S., see Ward, L.C., 353
Matsushita, M. and Wang, C.L., Projection pattern of vesitbulocerebellar fibers in the anterior vermis of the cat: an anterograde wheat germ agglutinin–horseradish peroxidase study, 25
Mediavilla, M.D., see Cos, S., 119
Meldrum, B.S., see Patel, S., 243
Menini, C., see Brailowsky, S., 75
Millan, M.H., see Patel, S., 243
Millar, T.J. and Morgan, I.G., Cholinergic amacrine cells in the rabbit retina synapse onto other cholinergic amacrine cells, 281
Molander, C., see Carlstedt, T., 14
Molander, C., Ygge, J. and Dalsgaard, C.-J., Substance P-, somatostatin- and calcitonin gene-related peptide-like immunoreactivity and fluoride resistant acid phosphatase-activity in relation to retrogradely labeled cutaneous, muscular and visceral primary sensory neurons in the rat, 37
Morgan, I.G., see Millar, T.J., 281
Morii, N., see Itoh, H., 102
Morreale, R.A., see Ferri, G.-L., 304
Morris, R., Responses of neurones in the brainstem raphe nuclei to stimulation of the cerebellar fastigial nuclei in the cat, 19

Nagatsu, T., see Naoi, M., 232
Nakao, K., see Itoh, H., 102
Naoi, M., Ishiki, R., Nomura, Y., Hasegawa, S. and Nagatsu, T., Quinolinic acid: an endogenous inhibitor specific for type B monoamine oxidase in human brain synaptosomes, 232
Naquet, R., see Brailowsky, S., 75
Nieuwenhuys, R., see Buma, P., 151
Nomura, Y., see Naoi, M., 232

Ohmori, Y., Watanabe, T. and Fujioka, T., Projections of visceral and somatic primary afferents to the sacral spinal cord of the domestic fowl revealed by transganglionic transport of horseradish peroxidase, 175
Oku, R., Satoh, M. and Takagi, H., Release of substance P from the spinal dorsal horn is enhanced in polyarthritic rats, 315
Oliver, C., see Ixart, G., 85

Palisses, R., see Viala, D., 49
Panerai, A.E., Sacerdote, P., Brini, A. and Mantegazza, P., Analgesic effect of morphine: a role for β-endorphin, 348
Patel, S., Millan, M.H. and Meldrum, B.S., Neurotransmission in the pendunculopontine nucleus and pilocarpine-induced motor limbic seizures in rats, 243
Pearson, J.C. and Goldfinger, M.D., The morphology and distribution of serotonin-like immunoreactive fibers in the cat dorsal column nuclei, 125
Pelletier, G., see Cailliez, D., 163
Penney, Jr., J.B., see Maragos, 371
Perry, T.L., Hansen, S. and Jones, K., Exposure to cigarette smoke does not decrease the neurotoxicity of *N*-methyl-4-phenyl-1-2, 3, 6-tetrahydropyridine in mice, 217
Persegol, L., see Viala, D., 49
Pin, J.-P., see Recasens, M., 211
Pinter, M.J., see Alstermark, B., 291

Polak, J.M., see Cailliez, D., 163
Post, C., see Fasmer, O.B., 81
Puelles, L., see Martinez-de-la-Torre, M., 31

Recasens, M., Pin, J.-P., and Bockaert, J., Chloride transport blockers inhibit the chloride-dependent glutamate binding to rat brain membranes, 211
Renaud, B., see Astier, B., 132
Reuss, S. and Schröder, H., Neuropeptide Y effects on pineal melatonin synthesis in the rat, 158
Risold, P.Y., see Fellmann, D., 275
Robitaille, R., Tremblay, J.P. and Grenon, G., Non-uniform distribution of miniature endplate potential amplitudes along the length of the frog neuromuscular junction, 187
Rodnight, R., see Dosemeci, A., 325
Roques, B.P., see Böhme, G.A., 227
Ross, C.D., Bowers, M. and Godfrey, D.A., Distributions of the activities of aspartate aminotransferase isoenzymes in rat retinal layers, 205
Roth, R.H., see Deutch, A.Y., 169
Rydqvist, B., see Kaila, K., 53
Rijnberk, A., see Stolp, R., 337

Sacerdote, P., see Panerai, A.E., 348
Saito, Y., see Itoh, H., 102
Salerno, M.T., see Cortimiglia, R., 193
Sánchez Barceló, E.J., see Cos, S., 119
Saper, C.B. and German, D.C., Hypothalamic pathology in Alzheimer's disease, 364
Satoh, M., see Oku, R., 315
Savasta, M., Dubois, A., Feuerstein, C., Manier, M. and Scatton, B., Denervation supersensitivity of striatal D_2 dopamine receptors is restricted to the ventro- and dorsolateral regions of the striatum, 180
Scatton, B., see Savasta, M., 180
Schröder, H., see Reuss, S., 158
Shiono, S., see Itoh, H., 102
Silva-Barrat, C., see Brailowsky, S., 75
Simon, W., see Ammann, D., 221
Slivka, A., Spina, M.B. and Cohen, G., Reduced and oxidized glutathione in human and monkey brain, 112
Soimero, L, see Ferri, G.-L., 304
Spina, M.B., see Slivka, A., 112
Steinbusch, H.W.M., see Stolp, R., 337
Stolp, R., Steinbusch, H.W.M., Rijnberk, A., and Croughs, R.J.M., Organization of ovine corticotropin-releasing factor immunoreactive neurons in the canine hypothalamo-pituitary system, 337
Stutzmann, J.-M., see Böhme, G.A., 227
Sugawara, A., see Itoh, H., 102
Swerup, C., see Kaila, K., 53

Takagi, H., see Oku, R., 315
Tamir, H., Liu, K.P. and Weiss, K.R., Proteins that bind serotonin in the nervous system of *Aplysia californica*, 257
Tantisira, B., see Alstermark, B., 286
Tantisira, B., see Alstermark, B., 291
Tiernan, J.M., see Ward, L.C., 353
Tork, I., see Bandler, R., 1
Tossman, U., see Lindefors, N., 90
Tremblay, J.P., see Robitaille, R., 187

Undén, A., see Westlind-Danielsson, A., 237
Ungerstedt, U., see Lindefors, N., 90

Valentino, R.J. and Aulisi, E.F., Carbachol-induced increases in locus coeruleus spontaneous activity are associated with an altered pattern of response to sensory stimuli, 297
Van Ree, J.M., see Király, I., 343
Vaudry, H., see Cailliez, D., 163
Viala, D., Persegol, L. and Palisses, R., Relationship between phrenic and hindlimb extensor activities during fictive locomotion, 49
Voipio, J., see Kaila, K., 53

Wainer, B.H., see Levey, A.I., 7
Wang, C.L., see Matsushita, M., 25
Ward, L.C., Tiernan, J.M., Hayling, N. and Mason, S., Protein turnover in subcellular fractions of brain from the ethanol-fed rat, 353
Watanabe, K., see Itoh, H., 102
Watanabe, T., see Ohmori, Y., 175
Weiss, K.R., see Tamir, H., 257
Westlind-Danielsson, A., Undén, A., Abens, J., Andell, S. and Bartfai, T., Neuropeptide Y receptors and the inhibition of adenylate cyclase in the human frontal and temporal cortex, 237
Wiesner, J.B. and Henriksen, S.J., Enkephalin enhances responsiveness to perforant path input while decreasing spontaneous activity in the dentate gyrus, 95

Yamada, T., see Itoh, H., 102
Yamaguchi, T., Monopodal fictive locomotion evoked by cervical cord stimulation in decerebrate cats, 69

Ygge, J., see Molander, C., 37
Young, A.B., see Maragos, W.F., 371

Zhang, L. and Krnjević, K., Apamin depresses selectively the after-hyperpolarization of cat spinal motoneurons, 58